

03-20-00

Attorney Docket No. 4600-0131.30

A

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE  
Certification under 37 CFR §1.10 (if applicable)

EL 530 369 535 US  
Express Mail Label Number

March 16, 2000  
Date of Deposit

I hereby certify that this Transmittal Letter, enclosed application and any other documents referred to as enclosed herein, are being deposited in an envelope with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR §1.10 on the date indicated above and addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231.

Matthew D. Redlon  
(Print Name of Person Mailing Application)

(Signature of Person Mailing Application)

**Transmittal of Utility Patent Application**  
**for Filing Under 37 CFR §1.53(b)**

Box Patent Application  
Assistant Commissioner for Patents  
Washington, D.C. 20231

Sir:

Transmitted herewith for filing is a utility patent application by inventors: Jagdish Parasrampur, Maxine B. Yonker, Kenneth E. Schwartz, Marc J. Gurwith, and entitled:

**DHEA COMPOSITION AND METHOD**

1. Enclosed are:

- ☒ This Transmittal letter.
- ☒ One stamped, self-addressed postcard for PTO date stamp.
- ☒ Certificate of Express Mail.
- ☒ One utility patent application containing text pages 1-23 and  
☒ 4 Sheets of drawings.
- ☒ Declaration of inventorship (unsigned)
- ☐ \* Assignment(s) for recordation with transmittal sheet.
- ☐ \* Small Entity Statement(s).
- ☐ Executed Power of Attorney by Assignee
- ☐ Sequence listing printout, diskette, and matching declaration.

2. U.S. Priority

- ☒ This application claims priority of Ser. No. 60/125,201 filed March 18, 1999, which is incorporated therein by reference.
- ☐ A petition for extension of time has been filed in the parent to extend the pendency of the parent to \* (copy enclosed).
- ☒ Conditional Petition for Extension of Time: An Extension of Time is requested to provide for timely filing if required to establish copendency with the parent after all papers filed herewith have been considered.

3. Foreign Priority

- ☐ Priority of Application No. \* filed in \* on \* is claimed under 35 USC §119.
- ☐ A certified copy of this priority document is enclosed.

4. Fees

The filing fee has been calculated as shown below:

For:	(Col. 1) No. Filed	(Col. 2) No. Extra	Small Entity		Other Than a Small Entity	
			Rate	Fee	Rate	Fee
Basic Fee				\$345.00	or	\$690.00
Total Claims	35 - 20	15	15 x \$ 9 =	\$135.00	or	x \$ 18 = \$
Independent Claims	7 - 3	4	4 x \$39 =	\$156.00	or	x \$ 78 = \$
<input type="checkbox"/> Multiple Dependent Claim Presented			+ \$130 =	\$ 00.00	or	+ \$260 = \$
* If the difference in Col. 1 is less than zero, enter "0" in Col. 2.						\$
TOTAL				\$636.00	or	TOTAL

- ☐ A check for \$\* is enclosed to cover the Filing Fee.
- ☐ Please charge Deposit Account 04-0531 in the amount of \$\*.
- ☐ The Commissioner is hereby authorized to charge fees under 37 CFR §1.16 and §1.17 which may be required, or credit any overpayment to Deposit Account 04-0531.
- ☒ Filing fee to be submitted in response to anticipated receipt of Notice to File Missing Parts of Application. DO NOT CHARGE DEPOSIT ACCOUNT.

Respectfully submitted,

*LeeAnn Gorthey*  
 LeeAnn Gorthey  
 Registration No. 37,337

Date: 3-16-2000

Correspondence Address:  
 Dehlinger & Associates  
 P.O. Box 60850  
 Palo Alto, CA 94306  
 (650) 324-0880  
 Customer No. 22918

**Transmittal of Utility Patent Application for Filing**

Certification Under 37 C.F.R. §1.10 (if applicable)

**EL 530 369 535 US**

"Express Mail" Label Number


**March 16, 2000**

Date of Deposit

I hereby certify that this Transmittal letter, enclosed application, and any other documents referred to as enclosed herein are being deposited in an envelope with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231

**Matthew D. Redlon**

(Print Name of Person Mailing Application)



(Signature of Person Mailing Application)

**DHEA Composition and Method**

This application claims priority to US provisional application serial number 60/125,201, filed March 18, 1999, which is hereby incorporated by reference in its entirety.

**Field of the Invention**

The present invention provides pharmaceutical formulations of DHEA enriched for polymorph form I or form II, which are useful for various therapeutic applications. In particular, the invention is directed to formulations of DHEA having more consistent bioavailability than previously used formulations.

**References**

Arlt, W. *et al.*, *J. Clin. Endocrinol. Metab.* **83**(6):1928-1934 (1998).

Barker, E.V. *et al.*, *Endocrinology* **134**:982-989 (1994).

Barrett-Connor *et al.*, *New Engl. J. Med.* **315**:1519 (1986).

Caira, M.R. *et al.*, *J. Chem. Crystallogr.* **25**:393 (1995).

Chang, L.C. *et al.*, *J. Pharmaceut. Sci.* **84**:1169-1179 (1995).

Comer, K.A. and Falany, C.N., *Mol. Pharmacol.* **41**:645-651 (1992).

Cox, P.J. *et al.*, *Acta Crystallogr.* **C46**, 334-336 (1990).

Falany, C.N. *et al.*, *Ann. NY Acad. Sci.* **774**:59-72 (1995).

Frye and Maciel, *J. Mag. Res.* **48**:125 (1982).

Grodin, J.M. *et al.*, *J. Clin. Endo. Metab.* **36**:207-214 (1973).

**GUIDANCE FOR INDUSTRY: Q2B VALIDATION OF ANALYTICAL PROCEDURES: METHODOLOGY**

HFD-210, CDER, Rockville, MD.

Kuhnert-Brandstätter, M., **THERMOMICROSCOPY IN THE ANALYSIS OF PHARMACEUTICALS**,

Pergamon Press, Oxford, U.K. (1971).

Lacheline, G.C. *et al.*, *J. Clin. Endocrinol. Metab.* **49**(6):892-898 (1979).

- Longcope, C., *Ann. NY Acad. Sci.* 774:143-148 (1995).
- MacDonald, J.C., *J. Mag. Res.* 38:381 (1980).
- Meikle, A.W. *et al.*, *J. Steroid Biochem. Molec. Biol.* 293-304 (1992).
- Orentreich, N. *et al.*, *J. Clin. Endocrinol. Metab.* 59:551-555 (1984).
- van Cauter, E., *Horm. Res.* 32(2):45-53 (1990).
- Yen, S.S. *et al.*, *Ann. NY Acad. Sci.* 774:128-142 (1995).

### **Background of the Invention**

Dehydroepiandrosterone (DHEA), also known as 3-beta-hydroxyandrost-5-en-17-one, dehydroisoandrosterone, trans-dehydroandrosterone,  $\Delta^5$ -androst-3- $\beta$ -ol-17-one, and prasterone, is a naturally occurring intermediate formed in the course of synthesis of various steroids from cholesterol. DHEA is the most abundant steroid hormone in humans and is produced mainly by the adrenal cortex as an inactive sulfate ester (DHEA-S). DHEA production also occurs in the testes, ovaries, and brain. After achieving a plateau level during early adulthood (ages 16 to 24), total serum DHEA (DHEA + DHEA-S) declines steadily to about 5 to 10% of peak values by age 60 to 70 (Orentreich *et al.*, 1984).

DHEA has been proposed for use in treating many medical conditions, such as systemic lupus erythematosus (U.S. Patent No. 5,817,650), primary adrenal insufficiency (U.S. Patent No. 5,861,391), Addison's disease (*ibid.*), reduced libido (U.S. Patent No. 5,855,548), obesity (U.S. Patent No. 5,846,962), osteoporosis (U.S. Patent Nos. 5,846,960 and 5,855,548), and fibromyalgia (U.S. Patent No. 5,935,949). DHEA can be administered by various routes and is orally active.

The pharmacokinetics of exogenously administered DHEA are complicated by endogenous production of DHEA and by the reversible interconversion between DHEA and DHEA-S, the major metabolite of DHEA, which acts as a reservoir for DHEA. DHEA exhibits wide diurnal variations in endogenous production, while DHEA-S levels show little variation during the day. Changes in plasma DHEA occur in parallel to those of ACTH and cortisol, with an early morning maximum, declining levels through the daytime, and minimal secretory activity in the early part of the night (van Cauter, 1990; Lacheline *et al.*, 1979; Yen *et al.*, 1995).

Both DHEA and DHEA-S are bound by serum albumin, globulins, and steroidal sex hormone binding globulin (Meikle *et al.*, 1992; Longcope, 1995). Only a small fraction of orally administered DHEA appears in the blood at any given time as DHEA; most undergoes conversion to DHEA-S by sulfotransferases in the liver and extrahepatic tissues (Barker, 1994; Comer, 1992; Falany, 1995; Arlt, 1998). DHEA-S is converted back to DHEA by peripheral tissues containing DHEA sulfatases, including lymphocytes and macrophages. DHEA is subsequently metabolized to androstenedione as well as the potent androgens, testosterone and dihydrotestosterone, and the estrogens, estrone and estradiol. Adipose tissue may serve as a substantial reservoir for adrenal androgens. The

aromatization of DHEA in peripheral tissue is thought to account for the majority of estrogen biosynthesis in postmenopausal women (Grodin *et al.*, 1973).

The bioavailability of a drug can play an important role in its efficacy. It has been reported that DHEA occurs in at least three and as many as five anhydrous polymorphic forms and at least three hydrated forms, depending on environmental conditions and the manner of preparation (Chang *et al.*, 1995). The known forms have been reported to be distinguishable on the basis of infrared spectroscopy and powder diffraction analysis, except that forms S3 and S4 are indistinguishable using the latter method (*ibid.*). Work conducted in support of the present invention indicates the existence of a sixth anhydrate form designated herein as form VI, which is detectable by solid state NMR but not by infrared spectroscopy or x-ray powder diffraction analysis.

Although DHEA is available from a variety of commercial sources, these materials show significant variation in their polymorphic compositions, which can cause variations in bioavailability due to differences of absorption during uptake *in vivo*.

Accordingly, it is an object of the present invention to provide DHEA formulations enriched in the form I polymorph or the form II polymorph, to achieve more consistent bioavailability and reliable efficacy. Formulations enriched in the form VI polymorph are also contemplated.

### **Summary of the Invention**

The present invention includes, in one aspect, a pharmaceutical formulation comprising dehydroepiandrosterone (DHEA), at least 85 % of which is present as the form I polymorph, and at least one pharmaceutical excipient. Preferably, at least 90 % of the DHEA is present as the form I polymorph, more preferably 95 %, and most preferably greater than 99 %. The invention also includes a composition of matter consisting essentially of the form I polymorph of DHEA.

The invention also includes, in a second general embodiment, a pharmaceutical formulation comprising DHEA, at least 85 % of which is present as the form II polymorph, and at least one pharmaceutical excipient. Preferably, at least 90 % of the DHEA is present as the form II polymorph, more preferably 95 %, and most preferably greater than 99 %.

Also included are pharmaceutical formulations comprising DHEA, at least 85 % of which is present as the form VI polymorph, and at least one pharmaceutical excipient. Preferably, at least 90 % of the DHEA is present as the form VI polymorph, more preferably 95 %, and most preferably greater than 99 %. The invention also includes a composition of matter consisting essentially of the form VI polymorph of DHEA.

The invention also includes a method for preparing a capsular or tablet formulation of DHEA. In the method, at least one solid pharmaceutical excipient is mixed with DHEA, at least 85 % of which is present as a single polymorph selected from form I and form II, and the solid formulation is either placed in a capsular container suitable for oral delivery or compressed to form a tablet.

In another aspect, the invention includes a method of administering DHEA to a subject to obtain an ameliorative result, wherein a pharmaceutically acceptable amount of DHEA is administered such that at least 85% of the DHEA is present as a single polymorph selected from form I, form II, and form VI, and preferably selected from forms I and II. These methods are useful for treating a variety of medical conditions, such as systemic lupus erythematosus, loss of bone density, osteoporosis, chronic fatigue syndrome, or fibromyalgia, or in DHEA replacement therapy.

The invention also includes a method for controlling the bioavailability of a DHEA formulation. In the method, a therapeutically effective amount of a DHEA formulation is administered to a subject, where the DHEA in the formulation consists of a preselected, known ratio of DHEA polymorphs.

The compositions and methods are useful for achieving more uniform bioavailabilities in DHEA formulations than have previously been achieved, in light of the applicants' discovery that bioavailability of DHEA *in vivo* is dependent upon the polymorphic composition of the DHEA.

These and other objects and features of the invention will be better understood in light of the following description.

### **Brief Description of the Drawings**

Figures 1 and 2 show mean baseline adjusted serum concentrations of DHEA and DHEA-S, respectively, in human subjects up to 72 hours after receiving a single dose of DHEA (formulation 1, 2 or 3, as described below) containing different proportions of DHEA forms I, II and VI; and

Figures 3 and 4 show mean baseline adjusted serum concentrations of DHEA and DHEA-S, respectively, in human subjects on days 7 to 10 of a multidose study, starting at 0.5 hours prior to dosing on day 7, using DHEA formulations, as described below, containing a mixture of forms I, II and VI (formulation 3) or pure form I (formulation 4).

### **Detailed Description of the Invention**

#### **I. DHEA Polymorphs**

The present invention is directed to pharmaceutical preparations of DHEA having more consistent bioavailabilities and pharmacokinetic properties than preparations available prior to the invention. In one aspect, the invention is directed to pharmaceutical compositions comprising DHEA such that at least 85%, preferably at least 90%, and more preferably at least 95% to 99% of the DHEA is present as the form I polymorph. Such compositions exhibit good uptake by the GI tract upon oral administration, show good therapeutic activity, and are highly stable under ambient conditions.

In another aspect, the invention is directed to pharmaceutical compositions comprising DHEA such that at least 85%, preferably at least 90%, and more preferably at least 95% to 99% of the DHEA is present as the form II polymorph. Such compositions exhibit good uptake by the GI tract

upon oral administration, a rapid rate of absorption (greater than the form I polymorph) and good therapeutic activity, and are also stable under ambient conditions.

Compositions thus enriched in form I or form II, as described herein, provide more predictable pharmacokinetic profiles than are provided by commercially available compositions having random polymorphic compositions. Compositions similarly enriched in form VI are also contemplated.

#### A. Preparation of DHEA Polymorphs

DHEA is known, via analytical techniques such as x-ray diffraction, infrared (IR) spectroscopy, and differential scanning calorimetry (DSC), to occur in several different hydrate and anhydrate crystal forms. The anhydrate forms include forms I, II, III, IV and V, although the latter two forms have been observed only transiently by DSC. The hydrates (solvates) include forms S1 (1/4 hydrate), S2 (monohydrate), S3 (monohydrate), and S4 (1/2 methanolate).

DHEA, and precursors such as DHEA acetate, are commercially available from various sources (e.g., Sigma Chemical Co., St. Louis, MO; Aldrich Chemical Company, Inc.; Diosynth, Inc.; Pfaltz & Bauer, Inc.; Schering AG). DHEA compositions enriched for selected polymorphs can be prepared by crystallization of commercial DHEA in selected solvents under appropriate cooling or evaporation conditions. Although conditions have been reported by others for preparing the above-mentioned stable forms (Chang *et al.*, 1995), it has been found by the present applicants that previously reported methods for preparing form I, including those described by Chang, yield products containing an additional polymorph that is designated herein as form VI. Accordingly, improved methods described herein have been developed for preparing pure form I to the exclusion of other polymorphs.

In one preferred method, pure form I is prepared by (a) crystallizing DHEA from anhydrous 2-propanol (or, alternatively, acetone or acetonitrile) under a nitrogen stream at room temperature over about 2 days, producing a crystalline precipitate that contains predominantly form I and some amount of form VI, followed by (b) suspending the precipitate in ethyl acetate (about 100 mL/30 g of DHEA) and stirring the resulting slurry at room temperature for about one week, followed by filtration. The filter cake is allowed to dry at room temperature overnight. <sup>13</sup>C-SSNMR analysis (discussed below) showed that product prepared by this method consisted of pure or nearly pure (>99%) form I; no other forms were detected by <sup>13</sup>C-SSNMR.

DHEA highly enriched for form II can be obtained by rapid crystallization from tetrahydrofuran (THF), dioxane, chloroform or mixtures of chloroform and THF. Example 1 provides a specific procedure for crystallization from THF, which produced a product shown by X-ray powder diffraction to be pure form II.

Other polymorphs can be prepared as follows. Form III can be obtained by desolvation of form S3 or S4 at room temperature under vacuum. Form VI can be obtained in pure form, or as a mixture

with form I, by crystallization of crude DHEA (prepared by saponification of DHEA acetate, as described in Example 1, without the methanol recrystallization step) from isopropanol, without a subsequent slurring step. Form S1 can be obtained by crystallization in methylene chloride under conditions of 50-60% relative humidity, or by grinding with methanol for 30 minutes and air drying.

Form S2 can be prepared by crystallization in 40% ethanol or distilled water, or by slow evaporation in acetonitrile, acetone, ethyl acetate, or THF. Form S3 can be prepared by replacing methanol molecules of a methanol half-solvate (form S4) with water under conditions of 60% relative humidity. Form S4 (1/2 methanolate) can be obtained by cooling a methanol solution of DHEA. Additional procedures for preparing various polymorphs can be found in Chang *et al.* (1995).

## B. Characterization of DHEA Compositions

In order to prepare DHEA compositions that are enriched for selected DHEA polymorphs, in accordance with the invention, it is important to be able to determine quantitatively the polymorph content of DHEA materials to establish the degree of enrichment for the desired polymorph relative to other polymorphs that may be present. Any appropriate method can be used for this purpose, provided that the method has a level of sensitivity and accuracy sufficient to permit measurement of the content of the major preferred polymorph within  $\pm 5\%$ , and preferably within  $\pm 2.5\%$  or less. Furthermore, it will be appreciated that a selected technique for measuring the polymorphs may not be capable of measuring all of the possible polymorphs, or it may be able to measure certain individual polymorphs only as a sum of two or more polymorphs. However, as discussed below, for measuring forms I, II and VI, these limitations either are not a concern or can be overcome by combining the results of complementary methods.

### B1. X-Ray Powder Diffraction

X-ray powder diffraction, the industry standard for determining polymorphic forms of crystalline substances, can be used to measure the relative amounts of forms I + VI, II, III, S1, S2, S3, and S4. However, this technique is not able to distinguish forms I and VI, as was shown by solid state nuclear magnetic resonance (see below).

In studies conducted in support of the present invention, essentially homogeneous samples of forms I, II, III, S1, S2, S3, and S4 of DHEA were prepared as described above. Form VI was studied as a 36:64 mixture (VI:I) with form I. Powder diffraction x-ray data were collected, as described in the Materials and Methods section below, to identify distinguishing peaks for each polymorph. The observed diffraction patterns were in general agreement with crystal structures reported previously for forms I, S1, S2, and S4 (Cox *et al.*, 1990; Caira *et al.*, 1995).

Table 1 summarizes the distinct reflections found for each polymorph. The relative amounts of each polymorph in a DHEA sample can be determined by integrating peak heights or peak areas of reflections that are unique to each DHEA polymorph in the sample, according to standard quantitative



x-ray methods, with integration of peak areas generally being preferred.

The diffractograms for the form I sample were found to be the same as for the form I/VI mixture, indicating that x-ray diffraction alone can quantify only the sum of these polymorphs. The level of one or the other form must be independently measured by another technique, preferably solid state NMR, as described below.

**Table 1. Unique Reflections of Polymorphs**

DHEA Form	Unique Peaks
I	15.0 (s), 16.8 (w), 18.0 (m), 18.7 (m), 19.1 (w), 19.3 (w), 20.2 (w), 24.8 (w), 25.0 (w), 25.2 (w)
II	8.6 (w), 17.3 (w), 20.9 (m), 22.0 (w), 22.2 (w), 27.1 (w)
III	11.8, 12.0, 13.0, 14.8, 17.0, 24.6
S1	13.4, 25.8
S2	11.1, 12.4, 14.1, 19.2
S3	14.1, 14.8, 18.2, 25.0
S4	14.0, 14.5, 17.1

<sup>a</sup> Peak positions are given in degrees 2 $\theta$ ; s = strong, m = medium, w = weak.

Typically, the XRPD analysis can be simplified by visually inspecting the diffractogram to determine which polymorphs are present at detectable levels, and then limiting integration to the unique reflections corresponding to those polymorphs. Usually, for anhydrous preparations, only forms I, II and/or VI are present, to the exclusion of hydrates S1 through S4. Minor polymorph components present at levels of about 5% can be readily quantified, with a lower limit of detection of about 2%.

For measuring polymorphic content of DHEA in pharmaceutical formulations containing additional materials such as pharmaceutical excipients, reflections arising from the additional materials can be subtracted out by calculating a difference pattern from diffractograms of such additional materials in the presence and absence of the DHEA.

## B2. Solid State NMR (SSNMR)

Solid state <sup>13</sup>C-NMR can also be used to detect and quantify polymorphs or crystalline substances and is a more sensitive technique than XRPD. However, as it is an expensive and time-consuming procedure, it is not routinely used. In the studies described herein, <sup>13</sup>C-SSNMR was required to distinguish forms I and VI of DHEA, as noted above.

Unique resonance peaks that can be used to quantify polymorphs I, II, VI, and S1-S4 are provided in Table 2. In general, the various polymorphs exhibit unique resonance shifts in the regions of 10-18 ppm (low field) and 115-124 ppm (high field) (ppm relative to adamantane). However, because some of the peaks overlap in the low field region, the high field region generally

provides better quantitative results.

Table 2. <sup>13</sup>C-SSNMR Peak Assignments

Form	Carbon No. 18 (ppm)	Carbon No. 6 (ppm)
In solution <sup>a</sup>	14.9	120.2
I	14.8, 14.1 <sup>b</sup>	120.4, 118.9 <sup>b</sup>
II	13.1	119.9
III	15.4, 13.1	120.1 <sup>c</sup>
VI	14.4	118.5
S1	14.3, 12.8 <sup>b</sup>	121.2, 119.5 <sup>b</sup>
S2	12.6 <sup>b</sup>	123.4, 120.4 <sup>b</sup>
S3	14.4	119.1
S4	13.9	119.1

5 <sup>a</sup>Pouchert, C.J., THE ALDRICH LIBRARY OF NMR SPECTRA, compound number 12,578-4 (1983).

<sup>b</sup>Known to have 2 crystallographically independent molecules in the structure.

The selected resonance peaks can be quantified by any known quantitation technique, such as the curve-fitting technique described by MacDonald (1980). Studies on mixtures of forms I and VI over a range of different concentrations showed that the responses to unique peaks for form VI are linear, indicating that the integration method is reliable for measuring the proportion of form VI in a sample relative to other polymorphs such as form I. Additional studies conducted in support of the invention established that there is good agreement for polymorph determinations by SSNMR and x-ray diffraction.

### B3. Infrared Spectroscopy

Infrared spectroscopy provides yet another method for determining polymorph content of DHEA compositions. Details of this technique can be found in Chang *et al.* (1995). However, like x-ray diffraction, it appears that this technique does not distinguish form I from form VI.

### C. Stability of DHEA Form I and Form II

A 14-day stress study monitored by XRPD showed that, at ambient temperature and 50°C, both form I and form II of DHEA are stable to solid form conversion for two weeks at relative humidities (RHs) as high as 84%. The 14-day data, given in Table 3, shows that Form I is stable for two weeks at ambient temperature and 95% RH. However, form II is converted to S2 under these conditions. At 50°C and 95% RH, form I is converted to S2; at 50° and at RH of 75% or greater, form II is partially converted to S1. Little weight gain (<0.8%) is observed in these samples. (Weight losses observed at 50°C could be due to loss of residual solvent, sublimation of sample, and/or thermal degradation.)

After one, two, three, and six months at either 25°C / 58% RH or 40°C / 75% RH, form I

remained stable to solid form conversion, as shown in the Table.

In summary, forms I and II are both stable under ambient conditions and under heat at non-saturating humidity; form I exhibits greater overall stability.

**Table 3. Stability of DHEA Forms I and II**

Polymorph	Temp	RH, %	Wt change, % (14 days)	Final Form (by XRPD), 14 days	Final Form (by <sup>13</sup> C-SSNMR), 1, 2, 3, and 6 months
Form I	25°C	58	-0.6	I	I
		75	+0.3	I	- (a)
		84	+0.1	I	-
		95	+0.2	I	-
	50°C	58	-1	I	-
		75	-3	I	I (b)
		84	-2	I	-
		95	+1	S2	-
Form II	25°C	58	0	II	
		75	-0.3	II	
		84	+0.2	II	
		95	+5	S2	
	50°C	58	-1	II	
		75	-2	II + S1	
		84	-0.3	II + S1	
		95	+1	II + S1	

(a) not determined

(b) temperature was 40°C.

## II. Formulation and Administration

Formulations containing the DHEA compositions of the invention may be provided in various dosage forms, such as, for example, tablets, capsules, powders, controlled-release formulations, suspensions, emulsions, suppositories, creams, ointments, lotions, or aerosols, and are preferably provided in solid dosage forms suitable for simple administration of precise dosages.

The compositions typically include a conventional pharmaceutical carrier, or excipient, and may additionally include other medicinal agents, carriers, adjuvants, and the like. Preferably, the composition will contain about 0.5% to 75% by weight, and more preferably about 5% to 25% by weight, DHEA, with the remainder consisting of suitable pharmaceutical excipients. For oral administration, such excipients may include pharmaceutical grades of lactose, mannitol, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, gelatin, sucrose, magnesium carbonate, and the like.

Conveniently, the formulations of the invention can be administered orally, e.g., by capsule or tablet, for rapid uptake in the blood stream and distribution to various compartments of the body. The amount and frequency of administration will vary, depending on the patient and on the therapeutic application as detailed further below. In another approach, solid formulations can be

administered as a suppository, e.g., when oral administration is contraindicated.

When the composition is employed in the form of solid preparations for oral administration, the preparations may be tablets, granules, powders, capsules or the like. For preparation of orally deliverable capsules, DHEA having the desired polymorph composition is mixed with at least one pharmaceutical excipient, and the solid formulation is placed in a capsular container suitable for delivery to the gastrointestinal tract. In preferred forms, at least 90%, more preferably at least 95%, and most preferably greater than 99% of the DHEA is present as a single polymorph selected from form I, II or VI, preferably form I or II, and most preferably form I.

For preparation of orally deliverable tablets, DHEA having the desired polymorph composition is mixed with at least one pharmaceutical excipient, and the solid formulation is compressed to form a tablet according to known methods, for delivery to the gastrointestinal tract. The tablet composition is typically formulated with additives, e.g. a saccharide or cellulose carrier, a binder such as starch paste or methyl cellulose, a filler, a disintegrator, or other additives typically usually used in the manufacture of medical preparations. The composition of the invention may also be administered to a subject transdermally or by inhalation. Methods for preparing various conventional dosage forms are known or will be apparent to those skilled in the art; for example, see *Remington's Pharmaceutical Sciences* (19th Ed., Williams & Wilkins, 1995).

### III. Pharmacokinetics

As the data below demonstrate, DHEA formulations enriched for polymorph form I or form II are both absorbed well *in vivo* and are therapeutically active. The applicants have further discovered that commercially available supplies of DHEA can vary significantly in their polymorphic compositions, potentially resulting in significant variations in therapeutic bioavailability and efficacy. These problems are overcome by the present invention, which provides DHEA formulations of known composition having more reliable therapeutic properties, particularly consistent bioavailability.

#### A. Single Dose Study

Example 2 describes a study in which 34 healthy postmenopausal women were administered orally one of three formulations of DHEA, each prepared from commercially available DHEA, in capsular form. The formulations were administered in single dosages of four opaque, size 2 capsules, each capsules containing 50 mg DHEA, 152 mg lactose (169 mg for formulation 3), 108 mg corn starch, and 3 mg magnesium stearate.

The polymorphic compositions of the formulations differed as summarized in Table 4 below. Formulation 1 contained 82% form I, 0% form II, and 18% form VI, while formulation 2 contained 22% form I, 44% form II, and 33% form VI (by SSNMR). Formulation 3, fairly similar to formulation 2, contained 18% form I, 43% form II, and 39% form VI. The average particle sizes in these formulations varied according to the technique used for measurement but generally fell within a

range of 50 - 150  $\mu\text{m}$ , with the particles in formulation 1 being somewhat larger (Table 5).

**Table 4. Polymorph Ratios in Formulations 1-3**

Formulation	[I + VI]:II Ratio (by XRPD)	I:VI:II Ratio (by SSNMR)
1	100 : 0	82 : 18 : 0
2	53 : 47	22 : 33 : 44
3	49 : 51	18 : 43 : 39

**Table 5. Particle Size**

Methodology	Formulation			
	1	2	3	4 (see below)
Microscopy	97 $\mu\text{m}$	55 $\mu\text{m}$	74 $\mu\text{m}$	-
Sonic Sifter Sieving	207 $\mu\text{m}$	180 $\mu\text{m}$	180 $\mu\text{m}$	-
Malvern Light Scattering	141 $\mu\text{m}$	73 $\mu\text{m}$	88 $\mu\text{m}$	71 $\mu\text{m}$

Blood samples were collected prior to and at various times after administration, and the levels of DHEA and DHEA-S were measured by immunoassay. Time plots of the levels of DHEA and DHEA-S are shown in Figs. 1 and 2, respectively. With reference to Fig. 1, the profile for formulation 1 shows that a maximum concentration was reached about three hours after administration, followed by a somewhat rapid decline to half the maximum concentration after about 12 hours and then a gradual decline to about a third of the maximum concentration after 72 hours. Formulations 2 and 3 achieved a maximum concentration more rapidly, within about an hour of administration, followed by a 2-stage decline in concentration similar to that shown by formulation 1.

These results show that both form I and form II have high bioavailability when administered orally. The data also indicate that formulations 2 and 3, having a high proportion of form II, are absorbed more rapidly and appear to have a higher bioavailability (by about 6-14% based on AUC measurements) relative to formulation 1.

The profiles for DHEA-S showed similar patterns. All formulations reached a maximum concentration after about three hours, showing that DHEA was converted rapidly to its major metabolite, the sulfate form. The shapes of the profiles were also very similar, except that formulation 1 showed consistently lower levels of DHEA-S than formulations 2 and 3, consistent with the results for DHEA shown in Fig. 1. The results show again that formulation 1 appears to have a lower bioavailability than formulations 2 and 3.

The estimated mean pharmacokinetic parameters (see Example 1) from this study are presented in Tables 6 and 7 below. AUC (area under curve) is an indicator of extent of absorption;  $C_{\text{max}}$  and  $T_{\text{max}}$  are indicators of rate of absorption. Serum DHEA concentrations for all samples were above the assay limit of quantification. Baseline-adjusted values were obtained by subtracting the day 1 pre-dosing serum concentration from each serum concentration value.

**Table 6: Baseline Adjusted Mean Pharmacokinetic Parameters for Serum DHEA: Single Dose Study**

Parameters	Mean (Std Dev); n=34		
	Formulation 1	Formulation 2	Formulation 3
$C_{\max}$ ( $\mu\text{g}/\text{dL}$ )	1.25 (0.614)	1.70 (1.07)	1.68 (0.918)
$T_{\max}$ (hr)	2.78 (0.915)	1.84 (1.16)	2.09 (1.34)
$AUC_{(0-\infty)}$ ( $\mu\text{g} \cdot \text{hr}/\text{dL}$ )	24.0 (9.89)	25.6 (10.2)	25.2 (8.53)
$AUC_{(0-72)}$ ( $\mu\text{g} \cdot \text{hr}/\text{dL}$ )	19.4 (8.56)	22.0 (8.43)	21.1 (7.31)

**Table 7: Baseline Adjusted Mean Pharmacokinetic Parameters for Serum DHEA-S: Single Dose Study**

Parameters	Mean (Std Dev); n=34		
	Formulation 1	Formulation 2	Formulation 3
$C_{\max}$ ( $\mu\text{g}/\text{dL}$ )	732.4 (193.0)	910.6 (251.7)	873.2 (230.1)
$T_{\max}$ (hr)	2.68 (0.72)	2.22 (0.97)	2.32 (0.70)
$AUC_{(0-\infty)}$ ( $\mu\text{g} \cdot \text{hr}/\text{dL}$ )	14396 (5591)	15851 (6461)	15290 (5818)
$AUC_{(0-72)}$ ( $\mu\text{g} \cdot \text{hr}/\text{dL}$ )	12395 (4969)	14423 (5569)	14121 (5175)

**B. Multiple Dose Study**

Example 3 describes an open-label, randomized, three period crossover pharmacokinetic study in which 39 healthy postmenopausal women were administered orally one of two formulations of DHEA in capsular form. The formulations were administered in single dosages of four opaque, size 2 capsules, each capsule containing 50 mg DHEA, 152 mg (Formulation 4) or 169 mg (Formulation 3) lactose, 108 mg corn starch, and 3 mg magnesium stearate. The DHEA polymorph composition of formulation 3 was as shown in Table 4 above (18% form I, 43% form II, and 39% form VI by SSNMR). Formulation 4 contained substantially pure (~100%) form I DHEA, prepared by methods described herein.

Dosages were administered at the same time each morning of each of two seven-day study periods, separated by a seven-day washout period. Serum samples were obtained five minutes prior to dosing on days 1 to 6, for measurement of trough levels of DHEA and DHEA-S. On day 7 of each study period, serum samples were obtained 30 minutes prior to dosing, then at various time intervals up to 72 hours post dosing, and the levels of DHEA and DHEA-S were measured by immunoassay, as described in Example 3.

Tables 8 and 9 show baseline-adjusted pharmacokinetic parameters  $AUC_{144-168}$  (extent of absorption),  $T_{\max}$  and  $C_{\max}$  (rate of absorption) for the two formulations. The  $AUC_{144-168}$  value represents a single dosing interval (day 7), as is standard for multidose studies.

**Table 8. Baseline Adjusted Mean Pharmacokinetic Parameters for Serum DHEA:  
Multiple Dose Study**

Parameters	Mean (SD); n=39	
	Formulation 3	Formulation 4
$C_{max}$ (µg/dL)	1.04 (0.44)	0.94 (0.66)
$T_{max}$ (hr)	2.3 (0.8)	2.5 (0.6)
$AUC_{144-168}$ (µg·hr/dL)	9.011 (3.99)	8.36 (3.67)

**Table 9. Baseline Adjusted Mean Pharmacokinetic Parameters for Serum DHEA-S:  
Multiple Dose Study**

Parameters	Mean (SD); n=39	
	Formulation 3	Formulation 4
$C_{max}$ (µg/dL)	1477 (390)	1295 (358)
$T_{max}$ (hr)	2.3 (0.7)	2.4 (0.5)
$AUC_{144-168}$ (µg·hr/dL)	15226 (5262)	13995 (4800)

Consistent with the results shown in the single-dose study above, formulation 4 (~100% form I) showed somewhat reduced bioavailability, in terms of rate and degree of absorption, compared to formulation 3 (mixture of polymorphs).

Time plots of the levels of DHEA and DHEA-S, from 0.5 hours before dosing on day 7 through day 10, are shown in Figs. 3 and 4, respectively. Again, the plots show that, while both formulations are absorbed rapidly, formulation 3, having a high proportion of form II, is absorbed more rapidly and appears to have a higher bioavailability than formulation 4, containing pure form I.

#### IV. Bioequivalence in DHEA Formulations

In one aspect, the invention is directed to methods of controlling bioavailability of DHEA formulations and achieving bioequivalence between different formulations. For example, different DHEA formulations containing DHEA and excipients in the same proportions may be prepared from different batches of DHEA and thus may exhibit different bioavailabilities *in vivo*. As discussed above, it has been found by the inventors that, because DHEA preparations can vary significantly in their polymorph compositions (1) from different suppliers, and (2) from batch to batch from the same supplier, pharmaceutical formulations prepared from commercially available supplies of DHEA can vary significantly in their *in vivo* bioavailabilities.

The invention also includes a method for controlling the bioavailability of a DHEA formulation. In the method, a therapeutically effective amount of a DHEA formulation is administered such that the formulation comprises a preselected, known ratio of DHEA polymorphs. That is, a DHEA formulation is prepared having known proportions (and amounts) of one or more DHEA polymorphs, e.g., by using appropriate crystallization conditions as discussed above, or by mixing appropriate amounts of the desired polymorphs. The polymorphic ratio can be selected to provide a desired bioavailability of the

DHEA, e.g., based on assessment of the bioavailability by methods such as described above and in Example 2. The method is useful for achieving more uniform and predictable bioavailabilities of DHEA formulations than have previously been achieved, in light of the applicants' discovery that bioavailability of DHEA *in vivo* is dependent upon the polymorphic composition of the DHEA.

#### V. Indications

DHEA is reported to be useful in treating many medical conditions. Exemplary conditions which can be treated include, for example, systemic lupus erythematosus (McGuire *et al.*, U.S. Patent No. 5,817,650), where a dose of about 25 to 500 mg/day is administered, optionally concomitantly with a glucocorticoid such as prednisone. Corticosteroids may also be used to supplement DHEA, administered at 0.25 to 2.0 mg/kg, in treatment of primary adrenal insufficiency or Addison's disease (S.S.C. Yen *et al.*, U.S. Patent No. 5,861,391). DHEA can also be used to enhance the immune response to viral infection (U.S. Patent No. 5,077,284), e.g. in treatment of immunocompromised subjects, such as AIDS patients, e.g. with dosages of 400 mg/day. Diabetes can be treated by administration of 120-480 mg/kg daily, for example, or in an amount equaling approximately 0.1% to 0.4% by weight of food intake, according to Coleman *et al.* (U.S. Patent No. 4,518,595).

Loss of bone density, as in osteoporosis or osteopenia, can also be treated with DHEA (see, for example, Labrie, U.S. Patent No. 5,776,923), with a typical dose being about 32 mg/kg daily. DHEA is also effective for treatment of chronic fatigue syndrome or fibromyalgia; see, for example, White, U.S. Patent No. 5,935,949.

Other conditions of interest and suggested dosages are summarized in the Table below. As always, optimum dosages, and frequency and duration of treatment, may vary among individual subjects and can be determined by methods known to persons skilled in the art.

**Table 10: Selected Conditions Treatable with DHEA**

Condition or Treatment	Recommended Dosage, daily	Reference
Inhibition of platelet aggregation (e.g. stroke, restenosis)	100 – 2000 mg	D.M. Eich <i>et al.</i> , U.S. Patent No. 5,110,810
Mild depression	15 – 150 mg	A.J. Morales <i>et al.</i> , U.S. Patent No. 5,407,927
Chronic renal failure	400-1600 mg	W. Walser, U.S. Patent No. 5,591,736
Ischemic injury (e.g. stroke, trauma)	2 – 50 mg/kg	R.A. Daynes <i>et al.</i> , U.S. Patent No. 5,635,496
Post-reperfusion injury	2 – 200 mg/kg	B.A. Araneo <i>et al.</i> , U.S. Patent No. 5,846,963
Reduced libido	5 – 50 mg/kg	F. Labrie, U.S. Patent No. 5,855,548
Pulmonary hypertension	2 – 50 mg/kg	B.A. Araneo <i>et al.</i> , U.S. Patent No. 5,753,640



Myotonic dystrophy	50 – 500 mg	N. Ohsawa <i>et al.</i> , U.S. Patent No. 5,834,451
Mast cell related allergic reactions, e.g. asthma	2 – 20 mg/kg	T. Dowell <i>et al.</i> , U.S. Patent No. 5,859,000
Arthritis	20 – 400 mg oral; 800 – 2000 mg topical	R.F. Peat, U.S. Patent No. 4,628,052
Lowering LDL cholesterol	100 – 1800 mg	J.E. Nestler <i>et al.</i> , U.S. Patent No. 4,920,115
Retroviral infections, especially HIV	100 – 600 mg	P.T. Prendergast, U.S. Patent No. 4,956,355

From the foregoing, it can be seen how the objects and features of the invention are met. The invention provides pharmaceutical formulations of DHEA that are enriched for polymorph form I, form II, or form VI, preferably form I or form II, and most preferably form I. These formulations provide consistent bioavailabilities, in contrast to previously utilized DHEA formulations containing varying amounts of multiple DHEA polymorphs. Formulations enriched in DHEA form I are highly stable and provide pharmaceutical compositions that retain a constant level of DHEA activity over long periods of time (e.g., more than 1 or 2 years). Accordingly, formulations enriched in form I have long shelf-lives, which is highly desirable in a pharmaceutical product. Formulations enriched in form II are stable, though somewhat less stable than form I formulations, and can have a faster rate of absorption (and higher efficacy) *in vivo* than form I formulations. Accordingly, formulations enriched in form II are also therapeutically beneficial.

The invention can be further understood in light of the following examples, which illustrate but are not intended in any way to limit the invention.

## EXAMPLES

### Materials and Methods

X-Ray Powder Diffraction (XRPD). XRPD analyses were carried out on a Siemens D-500 X-ray Powder Diffractometer-Kristaloflex or a Shimadzu XRD-6000 X-ray powder diffractometer using Cu K $\alpha$  radiation (1.5406 Å).

The Siemens instrument was equipped with an IBM-compatible interface and utilized DIFFRAC AT software (SOCABIM, 1994). Slits I and II were set at 1° and the radiation was electronically filtered by a Kevex Psi Peltier cooled silicon detector with slits III at 1° and IV at 0.15°. A continuous theta-two theta scan at 6°/min (0.4 sec/0.04° step) from 4 to 40 °2 $\theta$  was used. A silicon standard was analyzed each day to check the instrument alignment. Each sample was prepared by pressing a small amount of the powder onto a zero-background quartz-in-aluminum sample holder.

The Shimadzu instrument was equipped with a fine-focus X-ray tube. The tube power was set at 40 kV, 40 mA. The divergence and scattering slits were set at 1° and the receiving slit was set at 0.15

mm. Diffracted radiation was detected by an NaI scintillation detector. A theta-two theta continuous scan at 3°/min (0.4 sec/0.02° step) from 4 to 40 °2θ was used. A silicon standard was analyzed each day to check the instrument alignment, and an alumina standard was analyzed each day to check the X-ray tube output. Each sample was prepared for analysis by pressing it with a spatula onto a glass or quartz sample holder.

Quantitative work (measurements of peak heights and areas) was generally done on the Shimadzu instrument. Measurements of XRPD peak heights and areas were carried out using GRAMS/32 version 5.05 running on Windows NT. For quantifying the ratio of FII to FI+FVI, theta-two theta continuous scans were performed over a selected range (e.g., from 17 to 23 °2θ) at a selected scan range of from 0.5 to 3.0 °/min, typically 1°/min (0.4 sec/0.02° step), three times for each sample. The XRPD files were converted to ASCII format and read into GRAMS/32 version 5.05 running on Windows NT. For each scan, the peak areas and heights of a unique form I peak (typically either 18.5 or 20.2 °2θ) and a unique form II peak (typically 20.8 °2θ as measured over 20.5-21.2 °2θ) were measured. A standard curve was generated using standardized mixtures of FI and FII ranging from 0 to 100% FII in FI at 10% intervals, except that a sample containing 5% FII was also included. The three heights (or areas) measured for each sample were averaged, and the averages were input into the equation defined by the calibration curve.

Solid State NMR Method (SSNMR). SSNMR spectra were obtained on a General Electric Omega PSG, 100 MHz spectrometer using about 50 mg of sample in a 5-mm diameter zirconia rotor. High-power proton decoupling and cross-polarization with magic-angle spinning at approximately 5 kHz were used. The magic angle was adjusted using the Br signal of KBr by detecting the side bands as described by Frye and Maciel, 1982. The chemical shifts were referenced externally to the CH resonance of adamantane at 29.50 ppm. Curve fitting of selected peaks for integration was performed as described by MacDonald (1980). The following peaks were used to measure forms I, II and VI: sum of peaks at 118.8 and 120.3 ppm (form I), peak at 119.8 ppm (form II), and peak at 118.5 ppm (form VI).

Determination of Polymorph Content. In order to determine the amounts of the various possible polymorphs of DHEA in DHEA-containing materials, the following 3-step approach was used.

1. Visual examination of an XRPD pattern to qualitatively confirm that only anhydrous forms are present.
2. Quantitative XRPD analyses to determine the amount of form FII present in a mixture of forms FI, FII, and/or FVI.
3. Quantitative SSNMR analyses to determine the amount of forms FI and FVI present.

#### Example 1: Preparation of DHEA and Polymorphs I and II

Synthesis of DHEA. DHEA was prepared from DHEA acetate (obtained from Diosynth, Chicago, IL or Berlichem, Montville, NJ) by saponification using potassium carbonate in methanol. The product

was dissolved in 6 parts methanol at reflux, and charcoal was added and removed by filtration. The methanol was evaporated until a volume of 3 parts remained, and the solution was cooled to 15°C, maintained at this temperature for 1 hour, and filtered. The wet product was refluxed with 8.5 parts of water to remove the methanol, filtered, and dried under vacuum at 90°C. The loss on drying

specification for the final product was  $\leq 0.5\%$ , and the specification for residual methanol was  $\leq 0.01\%$ .

Preparation of Form I. Thirty grams of DHEA, as prepared above, were placed in a 500 mL flask under a nitrogen atmosphere. Anhydrous 2-propanol (isopropanol) was added until all the DHEA dissolved. The resulting solution was stirred under a nitrogen flow for 2 days, at which time all of the solvent had evaporated. SSNMR analysis indicated that the product contained a mixture of forms I and VI, with form I predominating. To convert the form VI component to form I, the product (about 30 g) was added to about 100 mL of ethyl acetate, and the resulting slurry was stirred at ambient temperature for 1 week and filtered. The filter cake was allowed to dry under ambient conditions overnight and then passed through a single sieve (75  $\mu\text{m}$ ), yielding 9.0 g of particles with sizes > 75  $\mu\text{m}$  and 3.2 g of particles with sizes < 75  $\mu\text{m}$ .  $^{13}\text{C}$ -SSNMR analysis of the fraction with particle size > 75  $\mu\text{m}$  showed that only form I was present.

Preparation of Form II. Thirty grams of DHEA were placed in a 500 mL flask under a nitrogen atmosphere. Anhydrous tetrahydrofuran was added until all the DHEA dissolved. The resulting solution was stirred under a nitrogen flow for 3 days, at which time all of the solvent had evaporated. The solid was removed from the flask and found by XRPD analysis to be solely form II (Siemens diffractometer).

#### Example 2: Pharmacokinetics of Orally Administered DHEA Formulations: Single Dose Study

This study was an open-label, randomized, three period crossover pharmacokinetic study in 34 healthy postmenopausal women. Subjects were contacted the evening prior to each dosing visit and reminded to begin an overnight fast, with nothing to eat or drink (no water permitted) for 10 hours prior to dosing. Serum samples for measurement of DHEA and DHEA-S were drawn thirty minutes prior to dosing and at 0, 0.5, 1, 1.5, 2, 3, 4, 6, 8, 10, 12, 16, 24, 48 and 72 hours after the subjects had received a 200 mg oral dose of DHEA (4 capsules of formulation 1 or 2) with 8 oz. of water. Each dosing period was followed by a 7 day washout period before the next administration.

DHEA levels were determined by radioimmunoassay (RIA) at Endocrine Sciences Inc., after non-polar solvent extraction. Method validation data demonstrated a recovery range of 92-99%, a limit of detection (LOD) of 18.9 ng/dL, a limit of quantitation (LOQ) of 87.5 ng/dL, intra-assay precision  $\leq 5\%$ , and inter-assay precision  $\leq 10\%$ . Internal assay controls were monitored via control charts. DHEA-S levels were determined by radioimmunoassay (RIA) at Endocrine Sciences Inc., after enzymolysis of DHEA-S. Method validation data demonstrated recovery range 86-106%, limit of detection (LOD) 3.6  $\mu\text{g/dL}$ , intra-assay precision  $\leq 7\%$ , and inter-assay precision  $\leq 10\%$ . Internal assay controls were monitored via control charts.

Each capsule contained 50 mg DHEA (formulation 1, 2 or 3) and excipients consisting of 152 mg lactose (169 mg for formulation 3), 108 mg corn starch, and 3 mg magnesium stearate. The DHEA polymorph contents of formulations 1-3 are listed in Table 4 in Section III above. The formulations were prepared in bulk amounts (e.g., 150 kg) by weighing and sieving all individual components through a size 14 stainless steel mesh screen to de-lump. All components except magnesium stearate were blended together for at least 10 minutes using a Patterson-Kelly Twin Shell 10 cubic ft dry V-blender. Magnesium stearate was added and blended into the mixture for at least 5 minutes. Homogeneity was verified by taking samples from different regions of the blender and quantifying the DHEA content on a weight basis by HPLC (isocratic, 1 mL/min 45:45:10 acetonitrile:water:methyl t-butylether, Synchropak column (C18 RP-P-100, 5 micron, 250 x 4.6 mm i.d. from Thinchrom, Inc., Lafayette, IN). After weighing, the blend was placed in opaque size 2 gelatin capsules (Capsugel or Shionogi) with 313 mg fill-weights.

Generation of Pharmacokinetic Data and Statistical Analysis: Microsoft® Excel, SAS® and WinNonlin® were used for dataset generation, statistical analysis, and pharmacokinetic (PK) evaluation of patient data.

Adjusted serum levels were calculated by subtracting baseline (average of -0.5 hr and 0 hr serum levels). Any adjusted value less than zero was set to zero. For DHEA and DHEA-S, the peak serum concentration ( $C_{max}$ ) and time to peak concentration ( $T_{max}$ ) were assessed from the concentration-time data. The maximal observed serum DHEA and DHEA-S concentration ( $C_{max}$ ) and the corresponding sampling times ( $T_{max}$ ) were determined for all treatments;  $T_{max}$  is expressed in hours following the commencement of each treatment.

DHEA and DHEA-S AUC (area under curve) values were determined by the linear trapezoidal method from hour 0 to hour 72 ( $AUC_{(0-72)}$ ) or the last measurable concentration. The area under the concentration time curve from zero to infinity ( $AUC_{0-\infty}$ ) was determined using the following equation:

$$AUC_{0-\infty} = AUC_{0-t} + C_{last}/k_{el}$$

where  $AUC_{0-t}$  is the area under the concentration-time curve from zero to the last measurable concentration ( $C_{last}$ ) as determined by the linear trapezoidal method and  $k_{el}$  is the terminal elimination rate constant. The terminal elimination rate constant was determined by using linear regression of the individual log concentrations versus time during the terminal elimination phase. The elimination rate constant was estimated by performing a regression of the natural logarithm of the concentrations on sampling time, for times in the specified range. Regressions were repeated first using the last three points, then the last four points. The half-life ( $t_{1/2}$ ) can be calculated by dividing 0.693 by  $k_{el}$ .

The main pharmacokinetic parameters used to describe DHEA and DHEA-S pharmacokinetics during treatments are the area under serum concentration curve from hour 0 to hour 72 ( $AUC_{(0-72)}$ ), the maximum serum concentration ( $C_{max}$ ), and the time to maximum serum concentration ( $T_{max}$ ). Comparisons of these parameters were made with an analysis of variance (ANOVA) model using the PROC GLM software (PC SAS, version 6.10).

### Example 3: Pharmacokinetics of Orally Administered DHEA Formulations: Multiple Dose Study

This study was an open-label, randomized, steady-state, two-treatment cross-over study of DHEA pharmacokinetics/pharmacodynamics in healthy postmenopausal females. DHEA was administered on each of 7 days of two study periods, with an intervening 7-day washout period (Days 1 to 7 and Days 15 to 21). Subjects received a single 200 mg oral dose (four 50 mg capsules) of DHEA at the same time each morning for the 7 days of each study period. Subjects were instructed to fast for 10 hours prior to each DHEA dose. Each capsule contained 50 mg of DHEA and pharmaceutical excipients (169 mg (Formulation 3) or 152 mg (Formulation 4) lactose, 108 mg corn starch, and 3 mg magnesium stearate) to result in a total capsule fill weight of 330 and 313 mg for formulations 3 and 4, respectively. Subjects were randomized to receive one of the two formulations during each study period. The DHEA polymorph composition of formulation 3 was as shown in Table 4 above (18% form I, 43% form II, and 39 % form VI). Formulation 4 contained substantially pure (~100%) form I DHEA, prepared by methods described herein.

Serum samples for the measurement of trough levels of DHEA and DHEA-S were obtained five minutes prior to the 200 mg oral dose of DHEA on days 1 to 6 of each study period.

On day 7 of each study period, each subject received the same breakfast (2 hours after dosing), lunch (6 hours after dosing) and dinner (10 hours after dosing). A full pharmacokinetic study was conducted, with serum samples for the measurement of DHEA and DHEA-S obtained 30 minutes prior to dosing and at 0, 1, 1.5, 2, 2.5, 3, 4, 6, 8, 12, 24, 36, 48, 60 and 72 hours following dosing.

DHEA levels were determined by radioimmunoassay (RIA) at Quest Diagnostics Inc. at Nichol's Institute. Method validation data demonstrated the lower limit of quantitation (LLQ) to be approximately 10 ng/dL. Pharmacokinetic data was generated and statistical analysis carried out essentially as described for Example 2, above. The estimated mean pharmacokinetic parameters for serum DHEA and DHEA-S concentrations from this study are summarized in Tables 8 and 9, above (Section IIIB).

While the invention has been described with reference to specific methods and embodiments, it will be appreciated that various modifications can be made without departing from the spirit of the invention.

It is claimed:

1. A pharmaceutical formulation comprising dehydroepiandrosterone (DHEA), at least 85% of which is present as the form I polymorph, and at least one pharmaceutical excipient.

2. The formulation of claim 1, wherein at least 90% of said dehydroepiandrosterone (DHEA) is present as the form I polymorph.

3. The formulation of claim 1, wherein at least 95% of said dehydroepiandrosterone (DHEA) is present as the form I polymorph.

4. The formulation of claim 1, wherein at least 99% of said dehydroepiandrosterone (DHEA) is present as the form I polymorph.

5. A method for preparing a solid DHEA formulation, said method comprising:  
mixing at least one solid pharmaceutical excipient with dehydroepiandrosterone (DHEA), at least 85% of which is present as the form I polymorph.

6. The method of claim 5, wherein at least 90% of said dehydroepiandrosterone (DHEA) is present as the form I polymorph.

7. The method of claim 5, wherein at least 95% of said dehydroepiandrosterone (DHEA) is present as the form I polymorph.

8. The method of claim 5, wherein at least 99% of said dehydroepiandrosterone (DHEA) is present as the form I polymorph.

9. The method of claim 5, further comprising the step of placing the solid formulation into a capsular container suitable for delivery to the gastrointestinal tract.

10. The method of claim 5, further comprising the step of compressing the solid formulation to form a tablet.

11. In a method for administering dehydroepiandrosterone (DHEA) to obtain an ameliorative result, the improvement comprising administering a pharmaceutically acceptable amount of DHEA, wherein at least 85% of the DHEA is present as the form I polymorph.

12. The method of claim 11, wherein at least 90% of the DHEA is present as the form I polymorph.

13. The method of claim 11, wherein at least 95% of the DHEA is present as the form I polymorph.

14. The method of claim 11, wherein at least 99% of the DHEA is present as the form I polymorph.

15. The method of claim 11, wherein said ameliorative result is treatment of systemic lupus erythematosus.

16. The method of claim 11, wherein said ameliorative result is prevention or reduction of loss of bone density.

17. The method of claim 11, wherein said ameliorative result is treatment of chronic fatigue syndrome or fibromyalgia.

18. A pharmaceutical formulation comprising dehydroepiandrosterone (DHEA), at least 85% of which is present as the form II polymorph, and at least one pharmaceutical excipient.

19. The formulation of claim 18, wherein at least 90% of said dehydroepiandrosterone (DHEA) is present as the form II polymorph.

20. The formulation of claim 18, wherein at least 95% of said dehydroepiandrosterone (DHEA) is present as the form II polymorph.

21. The formulation of claim 18, wherein at least 99% of said dehydroepiandrosterone (DHEA) is present as the form II polymorph.

22. A method for preparing a solid DHEA formulation, said method comprising:  
mixing at least one solid pharmaceutical excipient with dehydroepiandrosterone (DHEA), at least 85% of which is present as the form II polymorph.

23. The method of claim 22, wherein at least 90% of said dehydroepiandrosterone (DHEA) is present as the form II polymorph.

24. The method of claim 22, wherein at least 95% of said dehydroepiandrosterone (DHEA) is present as the form II polymorph.

25. The method of claim 22, wherein at least 99% of said dehydroepiandrosterone (DHEA) is present as the form II polymorph.

26. The method of claim 22, further comprising the step of placing the solid formulation into a capsular container suitable for delivery to the gastrointestinal tract.

27. The method of claim 22, further comprising the step of compressing the solid formulation to form a tablet.

28. In a method for administering dehydroepiandrosterone (DHEA) to obtain an ameliorative result, the improvement comprising administering a pharmaceutically acceptable amount of DHEA, wherein at least 85% of the DHEA is present as the form II polymorph.

29. The method of claim 28, wherein at least 90% of the DHEA is present as the form II polymorph.

30. The method of claim 28, wherein at least 95% of the DHEA is present as the form II polymorph.

31. The method of claim 28, wherein at least 99% of the DHEA is present as the form II polymorph.

32. The method of claim 28, wherein said ameliorative result is treatment of systemic lupus erythematosus.

33. The method of claim 28, wherein said ameliorative result is prevention or reduction of loss of bone density.

34. The method of claim 28, wherein said ameliorative result is treatment of chronic fatigue syndrome or fibromyalgia.

35. A method for controlling the bioavailability of a DHEA formulation, the method comprising: administering to a subject a DHEA formulation comprising DHEA and a pharmaceutical excipient, wherein said DHEA in said formulation consists of a preselected, known ratio of DHEA polymorphs.



ABSTRACT

Disclosed are improved pharmaceutical formulations comprising dehydroepiandrosterone (DHEA), enriched in selected polymorphic forms, for therapeutic applications. In one embodiment, the formulation comprises, in solid form, DHEA, at least 85% of which is present as a single polymorph selected from the form I polymorph or the form II polymorph, and at least one pharmaceutical excipient. Methods for making and using such compositions are also disclosed.

5

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

Fig. 1  
Mean Baseline Adjusted DHEA Concentrations




Fig. 2

## Mean Baseline Adjusted DHEA-S Concentrations




Fig. 3





Fig. 4



**INVENTORSHIP DECLARATION BY JOINT INVENTORS**

I HEREBY DECLARE THAT:

My residence, post office address, and citizenship are stated next to my name in PART A on hereof.

I believe I am the original, first, and joint inventor of the subject matter which is claimed and for which a patent is sought on the invention entitled

**DHEA COMPOSITION AND METHOD**

the specification of which:

- ☒ is attached hereto.
- ☐ was filed on \_\_\_\_\_ as Application Serial No. \_\_\_\_\_ and was amended on \_\_\_\_\_ (if applicable).

I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, §1.56.

I hereby claim priority benefits under Title 35, United States Code §119 of any foreign application(s) for patent or inventor's certificate listed in PART B hereof and have also identified in PART B on hereof any foreign application for patent or inventor's certificate having a filing date before that of the application of which priority is claimed.

I hereby claim the benefit under Title 35, United States Code §119(e) and §120, of any United States application(s) listed in PART C hereof and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of the first paragraph of Title 35, United States Code §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations §1.56 which occurred between the filing date of the prior application and any national or PCT international filing date of this application.

I hereby declare that all statements made herein of my knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Please direct all telephone calls to LeeAnn Gorthey at (650) 324-0880. Address all correspondence to:

**Dehlinger & Associates**  
**P.O. Box 60850**  
**Palo Alto, CA 94306-0850**

INVENTORSHIP DECLARATION BY JOINT INVENTORS

**PART A: INVENTOR INFORMATION AND SIGNATURE**

Full name of **FIRST** inventor: Jagdish Parasrampur

Citizenship: USA Post Office Address: 4012 Riddle Court  
Bridgewater, New Jersey 08807

Residence (if different):

Inventor's Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Full name of **SECOND** inventor: Maxine B. Yonker

Citizenship: GB Post Office Address: 427 Lakeview Way  
Emerald Hills, California 94062

Residence (if different):

Inventor's Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Full name of **THIRD** inventor: Kenneth E. Schwartz

Citizenship: USA Post Office Address: 511 Virginia Avenue  
San Mateo, California 94402

Residence (if different):

Inventor's Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Full name of **FOURTH** inventor: Marc J. Gurwith

Citizenship: USA Post Office Address: 13157 La Paloma Road  
Los Altos, California 94022

Residence (if different):

Inventor's Signature: \_\_\_\_\_ Date: \_\_\_\_\_

INVENTORSHIP DECLARATION BY JOINT INVENTORSPART B: CLAIM TO PRIORITY OF FOREIGN APPLICATION(S) UNDER 35 U.S.C. 119(a-d)

Country

App. No.

Filing Date

PART C: CLAIM TO PRIORITY OF U.S. PROVISIONAL AND NONPROVISIONAL APPLICATION(S) UNDER 35 U.S.C. 119(e) AND 120

Serial No.

Filing Date

Status

60/125,20118 March 1999☐

Patented

☒

Pending

☐

Abandoned

☐

Patented

☐

Pending

☐

Abandoned